

- 3 Betablokátory**  
prof. MUDr. Jindřich Špinar, CSc., FESC Interní kardiologická klinika, LF MU a FN, Brno  
prof. MUDr. Jiří Vítovec, CSc., FESC Interní kardiologická klinika, LF MU a FN u sv. Anny, Brno
- 3 Co je nového v léčbě dyslipidemií**  
prof. MUDr. Vladimír Soška, CSc. Oddělení klinické biochemie, FN u sv. Anny v Brně; International Clinical Research Center, FN u sv. Anny v Brně; Katedra laboratorních metod, II. interní klinika, LF MU, Brno  
MUDr. Ondřej Kyselák Oddělení klinické biochemie, FN u sv. Anny v Brně
- 4 Lze užívat atorvastatin ráno v rámci kombinované tablety Lipertance?**  
prof. MUDr. Hana Rosolová, DrSc. Centrum preventivní kardiologie, 2. interní klinika FN a LF UK, Plzeň
- 4 Léčba rezistentní hypertenze v roce 2016**  
doc. MUDr. Jan Václavík, Ph.D. I. interní klinika – kardiologická FN, Olomouc, a LF Univerzity Palackého, Olomouc
- 5 Aspirin a clopidogrel – klasický základ protidestičkové léčby**  
prof. MUDr. Jan Vojáček, DrSc., FESC, FACC I. interní kardiologická klinika LF UK a FN, Hradec Králové
- 5 Evolokumab v léčbě dyslipidemií se zaměřením na familiární hypercholesterolemii**  
doc. MUDr. Michal Vrablík, Ph.D. Centrum preventivní kardiologie, III. interní klinika 1. LF UK a VFN, Praha
- 5 Novinky v antihypertenzní léčbě**  
MUDr. Ivan Řiháček, Ph.D. | prof. MUDr. Miroslav Souček, CSc. II. interní klinika MU a FN u sv. Anny, Brno
- 6 Novinky v akutní léčbě mozkového infarktu**  
MUDr. Petr Polidar Neurologické oddělení Thomayerovy nemocnice, Praha  
doc. MUDr. Robert Mikulík, Ph.D. I. neurologická klinika LF MU a FN u sv. Anny, Brno
- 6 Pacient s reziduální anginou pectoris po PCI – kazuistika**  
MUDr. Jaroslav Brotánek Interní oddělení Thomayerovy nemocnice, Praha
- 6 Inhibitory ACE versus sartany v léčbě primární hypertenze**  
MUDr. Jiří Slíva, Ph.D. Ústav farmakologie 3. LF UK, Praha
- 7 Chronická pankreatitida – stálá pankreatologická výzva**  
prof. MUDr. Petr Dítě, DrSc. | MUDr. Martina Bojková Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě MUDr. Bohuslav Kianička, Ph.D. | MUDr. Tomáš Kupka II. interní klinika FN u sv. Anny, LF MU, Brno  
MUDr. Kateřina Kapounková, Ph.D. Katedra podpory zdraví, Fakulta sportovních studií MU, Brno  
doc. MUDr. Jana Dvořáčková, Ph.D. Ústav patologie FN, Ostrava Mgr. Jarmila Šímová | doc. MUDr. Arnošt Martínek, CSc. CGB laboratoř, Výzkumný institut AGEL, Ostrava prof. MUDr. Miroslav Souček, CSc. II. interní klinika FN u sv. Anny, LF MU, Brno
- 7 Pokroky v diagnostice a terapii akutní pankreatitidy**  
prof. MUDr. Petr Dítě, DrSc. | MUDr. Martina Bojková Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě Jarmila Šímová CGB laboratoř, a. s., Ostrava, člen skupiny Agel MUDr. Bohuslav Kianička, Ph. D. II. interní klinika FN u sv. Anny, LF MU, Brno MUDr. Tomáš Kupka | Mgr. Lenka Dovrtělová, Ph.D. | doc. MUDr. Arnošt Martínek, CSc. Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě  
prof. MUDr. Miroslav Souček, CSc. II. interní klinika FN u sv. Anny, LF MU, Brno
- 8 Diagnostika a příznaky Fabryho choroby**  
MUDr. Lubor Goláš II. interní klinika kardiologie a angiologie 1. LF UK a VFN, Praha
- 8 Léčba CHOPN – současnost a novinky**  
doc. MUDr. Norbert Pauk, Ph.D. Klinika pneumologie a hrudní chirurgie Nemocnice Na Bulovce a 3. LF UK, Praha
- 8 Léčba astmatu – současnost a novinky**  
doc. MUDr. Norbert Pauk, Ph.D. Klinika pneumologie a hrudní chirurgie Nemocnice Na Bulovce a 3. LF UK, Praha
- 8 Novinky v léčbě metabolického syndromu**  
prof. MUDr. Štěpán Svačina, DrSc., MBA III. interní klinika 1. LF UK a VFN, Praha
- 9 Fixní kombinace bazálního inzulínového analogu a GLP-1 po selhání bazálního inzulínu**  
prof. MUDr. Zdeněk Rušavý, Ph.D. | MUDr. Michal Žourek I. interní klinika FN a LF UK, Plzeň
- 9 Novinky v léčbě diabetika 2. typu**  
prof. MUDr. Jan Škrha, DrSc., MBA III. interní klinika – klinika endokrinologie a metabolismu, 1. LF UK a VFN, Praha

- 9 **DM2T: léčebný „switch“ pomocí dapagliflozinu, snížení hmotnosti jako přidaná hodnota – kazuistika**  
MUDr. David Bartoš Diabetologická ambulance, Centrum zdraví Sagena, Frýdek-Místek  
Diabetologická ambulance, Vítkovická nemocnice, a. s.
- 10 **Současné možnosti léčby obezity a zdravotně závažné nadváhy**  
MUDr. Petr Sucharda, CSc. III. interní klinika VFN a 1. LF UK, Praha
- 10 **Časná léčba inzulinem u pacientů s diabetes mellitus 2. typu**  
MUDr. Milan Flekač, Ph.D. | MUDr. Jan Šoupal III. interní klinika VFN a 1. LF UK, Praha

# Betablokátory

prof. MUDr. Jindřich Špinar, CSc., FESC Interní kardiologická klinika, LF MU a FN, Brno  
prof. MUDr. Jiří Vítovec, CSc., FESC Interní kardiologická klinika, LF MU a FN u sv. Anny, Brno

- 1 Bangalore, S. – Nessedli, F. H. – Kostis, J. B., et al.: Cardiovascular protection using beta-blockers: a critical review of the evidence. *J Am Coll Cardiol*, 2007, 50, s. 563–572.
- 2 Bangalore, S. – Bhatt, D. L. – Steg, P. G., et al.:  $\beta$ -blockers and cardiovascular events in patients with and without myocardial infarction. Post hoc analysis from the CHARISMA trial. *Circ Cardiovasc Qual Outcomes*, 2014, 7, s. 872–881.
- 3 Bangalore, S. – Makani, H. – Radford, M., et al.: Clinical outcomes with  $\beta$ -blockers for myocardial infarction: a meta-analysis of randomized trials. *Am J Med*, 2014, 127, s. 939–953.
- 4 Bradley, H. A. – Wiyongse, C. S. – Vilmino, J. A., et al.: How strong is the evidence for use of beta-blockers as first-line therapy for hypertension? Systematic review and meta-analysis. *J Hypertens*, 2006, 24, s. 2131–2141.
- 5 Dubrava, J.: Prieskum „3P při srdcovom zlyhavaní“ so zameraním na srdcovú frekvenciu. *Vnitřní Léč*, 2016, 62, s. 17–24.
- 6 Filipovský, J. – Widimský, J. jr. – Ceral, J., et al.: Diagnostické a léčebné postupy u arteriální hypertenze – verze 2012. Doporučení České společnosti pro hypertenzi. *Vnitřní Léč*, 2012, 58, s. 785–802.
- 7 Fleischmann, K. E. – Beckman, J. A. – Bulle, C. E., et al.: 2009 ACCF/AHA focused update on perioperative beta-blockade: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*, 2009, 54, s. 2102–2128.
- 8 James, P. A. – Oparil, S. – Carter, L., et al.: Evidence-based guideline for the management of high blood pressure in adults report from the panel members appointed to the Eighth Joint National Committee. 2014 (JNC 8). *JAMA*, doi:10.1001/jama.2013.284427.
- 9 Kaplan, N. M.: Beta-blockers in hypertension adding insult to injury. *J Am Coll Cardiol*, 2008, 52, s. 1490–1491.
- 10 Lindholm, L. H. – Carlberg, B. – Samuelsson, O.: Should betablockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*, 2005, 366, s. 1545–1553.
- 11 Misumi, N. – Harjai, K. – Kernis, S., et al.: Does oral beta-blocker therapy improve long-term survival in ST-segment elevation myocardial infarction with preserved systolic function? A meta-analysis. *J Cardiovasc Pharmacol Ther*, 10, 4, 2016, pii: 1074248415608011.
- 12 Seronde, M. F. – Geha, R. – Puymirat, E., et al.: Discharge heart rate and mortality after acute myocardial infarction. *Am J Med*, 2014, 127, s. 954–962.
- 13 Špác, J.: Jsou betablokátory stále moderní léky? *Acta medicae*, 2016, 2, s. 6–9.
- 14 The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) 2016 European Guidelines on cardiovascular disease prevention in clinical practice. *European Heart Journal*, 2016, doi:10.1093/eurheartj/ehw106.
- 15 The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC): 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *European Heart Journal*, 2016, doi:10.1093/eurheartj/ehw128.
- 16 UK National Institute for Health and Clinical Excellence and British Hypertension Society. *Hypertension: management of hypertension in adults in primary care*. 2006, NICE guidelines.
- 17 Vítovec, J. – Špinar, J.: Diuretika a betablokátory v léčbě hypertenze. *Interní medicína pro praxi*, 2012, 14, s. 458–460.
- 18 Vítovec, J. – Špinar, J.: Betapres – nová fixní kombinace pro preventivní kardiologii. *Hypertenze a kardiologická prevence*, 2014, 3, s. 30–32.
- 19 Vítovec, J. – Špinar, J.: Beta-blockers in the treatment of chronic heart failure. How should results of clinical studies be introduced into clinical practice. *Vnitřní Léč*, 2000, 46, s. 161–165.
- 20 Widimský, P. – Kala, P. – Rokyta, R.: Souhrn Doporučených postupů ESC pro diagnostiku a léčbu pacientů s akutním infarktem myokardu s elevací úseků ST z roku 2012. Připraven Českou kardiologickou společností. *Cor Vasa*, 2012, 54, s. 447–463.
- 21 Widimský, P. – Rokyta, R. – Hlinomaz, O.: Souhrn Doporučených postupů ESC pro diagnostiku a léčbu pacientů s akutním infarktem myokardu s elevací úseků ST z roku 2016. Připraven Českou kardiologickou společností. *Cor Vasa*, 2016, 58, s. e4–e28.

# Co je nového v léčbě dyslipidemií

prof. MUDr. Vladimír Soška, CSc. Oddělení klinické biochemie, FN u sv. Anny v Brně;  
International Clinical Research Center, FN u sv. Anny v Brně;  
Katedra laboratorních metod, II. interní klinika, LF MU, Brno  
MUDr. Ondřej Kyselák Oddělení klinické biochemie, FN u sv. Anny v Brně

- 1 Cuchel, M. – Bruckert, E. – Ginsberg, H. N., et al.: Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. *Eur Heart J*, 2014, 35, s. 2146–2157.
- 2 Baigent, C. – Blackwell, L. – Emberson, J., et al.: Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170 000 participants in 26 randomised trials. *Lancet*, 2010, 376, s. 1670–1681.
- 3 Cholesterol Treatment Trialists' (CTT) Collaboration, Baigent, C. – Blackwell, L., et al.: Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet*, 2010, 376, s. 1670–1681.
- 4 Vavřková, H. – Soška, V. – Rosolová, H., et al.: Czech Atherosclerosis Society Guidelines for the diagnosis and treatment of dyslipidemia in adults. *Vnitřní Léč*, 2007, 53, s. 181–187.
- 5 Soska, V. – Vavřková, H. – Vrablík, M., et al.: Opinion of the Czech Atherosclerosis Society's committee (CSAT) on the ESC/EAS guidelines related to the diagnostics and treatment of dyslipidemia issued in 2011. *Vnitřní Léč*, 2013, 59, s. 120–126.
- 6 Catapano, A. L. – Reiner, Z. – De Backer, G., et al.: ESC/EAS Guidelines for the management of dyslipidaemias The Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Atherosclerosis*, 2011, 217, s. 3–46.
- 7 Piepoli, M. F. – Hoes, A. W. – Agewall, S., et al.: 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts): Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J*, 2016.
- 8 Perk, J. – De Backer, G. – Gohlke, H., et al.: European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). *Eur Heart J*, 2012, 33, s. 1635–1701.
- 9 Canon, C. P. – Blazing, M. A. – Giugliano, R. P., et al.: Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*, 2015, 372, s. 2387–2397.
- 10 Scotty, R. – O'Brien, R. – Fulcher, G., et al.: Effects of fenofibrate treatment on cardiovascular disease risk in 9,795 individuals with type 2 diabetes and various components of the metabolic syndrome: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. *Diabetes Care*, 2009, 32, s. 493–498.
- 11 Jun, M. – Foote, C. – Lv, J., et al.: Effects of fibrates on cardiovascular outcomes: a systematic review and meta-analysis. *Lancet*, 2010, 375, s. 1875–1884.
- 12 Rajamani, K. – Colman, P. G. – Li, L. P., et al.: Effect of fenofibrate on amputation events in people with type 2 diabetes mellitus (FIELD study): a prespecified analysis of a randomised controlled trial. *Lancet*, 2009, 373, s. 1780–1788.
- 13 Cheb, E. Y. – Davis, M. D. – Danis, R. P., et al.: The effects of medical management on the progression of diabetic retinopathy in persons with type 2 diabetes: the Action to Control Cardiovascular Risk in Diabetes (ACCORD) Eye Study. *Ophthalmology*, 2014, 121, s. 2443–2451.
- 14 Alan, Z. – Baass, A. – Genest, J.: Proprotein convertase subtilisin/kexin type 9 (PCSK9): lessons learned from patients with hypercholesterolemia. *Clin Chem*, 2014, 60, s. 1380–1389.
- 15 Koren, M. J. – Giugliano, R. P. – Raal, F. J., et al.: Efficacy and safety of longer-term administration of evolocumab (AMG 145) in patients with hypercholesterolemia: 52-week results from the Open-Label Study of Long-Term Evaluation Against LDL-C (OSLER) randomized trial. *Circulation*, 2014, 129, s. 234–243.
- 16 Li, C. – Lin, L. – Zhang, W., et al.: Efficiency and safety of proprotein convertase subtilisin/kexin 9 monoclonal antibody on hypercholesterolemia: a meta-analysis of 20 randomized controlled trials. *J Am Heart Assoc*, 2015, 4, s. e001937.
- 17 Canon, C. P. – Cariou, B. – Blom, D., et al.: Efficacy and safety of alirocumab in high cardiovascular risk patients with inadequately controlled hypercholesterolemia on maximally tolerated doses of statins: the ODYSSEY COMBO II randomized controlled trial. *Eur Heart J*, 2015, 36, s. 1186–1194.
- 18 Raal, F. J. – Giugliano, R. P. – Sabatine, M. S., et al.: Reduction in lipoprotein(a) with PCSK9 monoclonal antibody evolocumab (AMG 145): a pooled analysis of more than 1,300 patients in 4 phase II trials. *J Am Coll Cardiol*, 2014, 63, s. 1278–1288.
- 19 Lipinski, M. J. – Benedetto, U. – Escarrega, R. O., et al.: The impact of proprotein convertase subtilisin-kexin type 9 serine protease inhibitors on lipid levels and outcomes in patients with primary hypercholesterolaemia: a network meta-analysis. *Eur Heart J*, 2016, 37, s. 536–545, doi: 10.1093/eurheartj/ehv563.
- 20 Robinson, J. G. – Farnier, M. – Krempf, M., et al.: Efficacy and safety of alirocumab in reducing lipids and cardiovascular events. *N Engl J Med*, 2015, 372, s. 1489–1499.
- 21 Sabatine, M. S. – Giugliano, R. P. – Wiviott, S. D., et al.: Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med*, 2015, 372, s. 1500–1509.
- 22 Navarese, E. P. – Kolodziejczak, M. – Schulze, V., et al.: Effects of proprotein convertase subtilisin/kexin type 9 antibodies in adults with hypercholesterolemia: A systematic review and meta-analysis. *Ann Intern Med*, 2015, 163, s. 40–51.
- 23 Ling, H. – Burns, T. L. – Hilleman, D. E.: An update on the clinical development of proprotein convertase subtilisin kexin 9 inhibitors, novel therapeutic agents for lowering low-density lipoprotein cholesterol. *Cardiovasc Ther*, 2014, 32, s. 82–88.
- 24 Raal, F. J. – Stein, E. A. – Divour, R., et al.: PCSK9 inhibition with evolocumab (AMG 145) in heterozygous familial hypercholesterolaemia (RUTHERFORD-2): a randomised, double-blind, placebo-controlled trial. *Lancet*, 2015, 385, s. 331–340.
- 25 Kastelein, J. J. – Ginsberg, H. N. – Langslet, G., et al.: ODYSSEY FH I and FH II: 78 week results with alirocumab treatment in 735 patients with heterozygous familial hypercholesterolaemia. *Eur Heart J*, 2015, 36, s. 2996–3003.
- 26 Stroes, E. – Colquhoun, D. – Sullivan, D., et al.: Anti-PCSK9 antibody effectively lowers cholesterol in patients with statin intolerance: the GAUSS-2 randomized, placebo-controlled phase 3 clinical trial of evolocumab. *J Am Coll Cardiol*, 2014, 63, s. 2541–2548.
- 27 Moriarty, P. M. – Jacobson, T. A. – Bruckert, E., et al.: Efficacy and safety of alirocumab, a monoclonal antibody to PCSK9, in statin-intolerant patients: design and rationale of ODYSSEY ALTERNATIVE, a randomized phase 3 trial. *J Clin Lipidol*, 2014, 8, s. 554–561.

- 28 Vuorio, A. – Tikkanen, M. J. – Kovanen, P. T.: Inhibition of hepatic microsomal triglyceride transfer protein – a novel therapeutic option for treatment of homozygous familial hypercholesterolemia. *Vasc Health Risk Manag*, 2014, 10, s. 263–270.
- 29 Voight, B. F. – Peloto, G. M. – Orho-Melander, M., et al.: Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet*, 2012, 380, s. 572–580.
- 30 Sarwar, N. – Danech, J. – Eiriksdottir, G., et al.: Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation*, 2007, 115, s. 450–458.
- 31 Langsted, A. – Freiberg, J. J. – Nordestgaard, B. G.: Fasting and non-fasting lipid levels: influence of normal food intake on lipids, lipoproteins, apolipoproteins, and cardiovascular risk prediction. *Circulation*, 2008, 118, s. 2047–2056.
- 32 Robinson, J. G. – Wang, S. – Smith, B. J., et al.: Meta-analysis of the relationship between non-high-density lipoprotein cholesterol reduction and coronary heart disease risk. *J Am Coll Cardiol*, 2009, 53, s. 316–322.

## Lze užívat atorvastatin ráno v rámci kombinované tablety Lipertance?

prof. MUDr. Hana Rosolová, DrSc.

Centrum preventivní kardiologie, 2. interní klinika FN a LF UK, Plzeň

- 1 *Zocor (simvastatin) package insert*. Whitehouse Station, NJ, Merck Pharmaceuticals, 2004.
- 2 *Mevacor (lovastatin) package insert*. Whitehouse Station, NJ, Merck Pharmaceuticals, 2004.
- 3 *Lescol (fluvastatin) package insert*. East Hanover, NJ, Novartis Pharmaceuticals, 2004.
- 4 *Pravachol (pravastatin) package insert*. Princeton, NJ, Bristol-Myers Squibb, 2003.
- 5 *Lipitor (atorvastatin calcium) package insert*. Morris Plains, NJ, Pfizer Pharmaceuticals, 2003.
- 6 *Crestor (rosuvastatin) package insert*. Wilmington, DE, AstraZeneca, 2003.
- 7 Plakogiannis, R. – Cohen, H.: Optimal LDL cholesterol lowering – morning versus evening statin administration. *Ann Pharmacother*, 2007, 41, s. 106–110.
- 8 Cilla, D. D. Jr. – Gibbon, D. M., et al.: Pharmacodynamic effects and pharmacokinetics of atorvastatin after administration to normocholesterolemic subjects in the morning and evening. *J Clin Pharmacol*, 1996, 36, s. 604–609.
- 9 Plakogiannis, R. – Cohen, H., et al.: Effects of morning versus evening administration of atorvastatin in patients with hyperlipidemia. *Am J Health-Syst Pharm*, 2005, 62, s. 2491–2494.
- 10 Kostis, W. J. – Cheby, J. Q. – Dobrzynski, J. M., et al.: Metaanalysis of statins effects in women versus men. *J Am Coll Cardiol*, 2012, 59, s. 572–582.
- 11 Chapman, R. H. – Benner, J. S. – Petrilla, A. A., et al.: Predictors of adherence with antihypertensive and lipid-lowering therapy. *Arch Intern Med*, 2005, 165, s. 1147–1152.

## Léčba rezistentní hypertenze v roce 2016

doc. MUDr. Jan Václavík, Ph.D.

I. interní klinika – kardiologická FN, Olomouc, a LF Univerzity Palackého, Olomouc

- 1 Calhoun, D. – Jones, D. – Textor, S., et al.: Resistant hypertension: diagnosis, evaluation, and treatment: a scientific statement from the American Heart Association Professional Education Committee of the Council for High Blood Pressure Research. *Circulation*, 2008, 117, s. e510–e526, doi:10.1161/CIRCULATIONAHA.108.189141.
- 2 Lancia, G. – Tabard, R. – Narkiewicz, K., et al.: 2013 ESH/ESC guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens*, 2013, 31, s. 1281–1357, doi:10.1093/eurheartj/ehs1151.
- 3 Persell, S. D.: Prevalence of resistant hypertension in the United States, 2003–2008. *Hypertension*, 2011, 57, s. 1076–1080, doi:10.1161/HYPERTENSIONAHA.111.170308.
- 4 Brambilla, G. – Bombelli, M. – Seravalle, G., et al.: Prevalence and clinical characteristics of patients with true resistant hypertension in central and Eastern Europe: data from the BP-CARE study. *J Hypertens*, 2013, 31, s. 2018–2024, doi:10.1097/HJH.0b013e328363823f.
- 5 Pierdomenico, S. D. – Lapena, D. – Bucci, A., et al.: Cardiovascular outcome in treated hypertensive patients with responder, masked, false resistant, and true resistant hypertension. *Am J Hypertens*, 2005, 18, s. 1422–1428, doi:10.1016/j.amjhyper.2005.05.014.
- 6 Neter, J. E. – Stam, B. E. – Kok, F. J., et al.: Influence of weight reduction on blood pressure: A meta-analysis of randomized controlled trials. *Hypertension*, 2003, 42, s. 878–884, doi:10.1161/01.HYP.0000094221.86888.AE.
- 7 He, F. J. – Li, J. – MacGregor, G. A.: Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ*, 2013, 346, f1325, dostupné z: <http://www.ncbi.nlm.nih.gov/pubmed/23558162>, vyhledáno 29. 5. 2015.
- 8 Dimeo, F. – Pagonas, N. – Seifert, F., et al.: Aerobic exercise reduces blood pressure in resistant hypertension. *Hypertension*, 2012, 60, s. 653–658, doi:10.1161/HYPERTENSIONAHA.112.197780.
- 9 Guimaraes, G. V. – de Barros Cruz, L. G. – Fernandes-Silva, M. M., et al.: Heated water-based exercise training reduces 24-hour ambulatory blood pressure levels in resistant hypertensive patients: A randomized controlled trial (HEX trial). *Int J Cardiol*, 2014, 172, s. 434–441, doi:10.1016/j.ijcard.2014.01.100.
- 10 Fontil, V. – Pletcher, M. J. – Khanna, R., et al.: Physician underutilization of effective medications for resistant hypertension at office visits in the United States: NAMCS 2006–2010. *J Gen Intern Med*, 2014, 29, s. 468–476, doi:10.1007/s11606-013-2683-y.
- 11 Sarafidis, P. – Bakris, G. L.: Resistant hypertension. An overview of evaluation and treatment. *J Am Coll Cardiol*, 2008, 52, s. 1749–1757, doi:10.1016/j.jacc.2008.08.036.
- 12 Finnerty, F. A. – Maxwell, M. H. – Lunn, J., et al.: Long-term effects of furosemide and hydrochlorothiazide in patients with essential hypertension a two-year comparison of efficacy and safety. *Angiology*, 1977, 28, s. 125–133, dostupné z: <http://www.ncbi.nlm.nih.gov/pubmed/869273>, vyhledáno 31. 5. 2015.
- 13 Williams, B. – MacDonald, T. M. – Morant, S., et al.: Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet*, 2015, 386, s. 2059–2068, doi:10.1016/S0140-6736(15)00257-3.
- 14 Hermida, R. C. – Ayala, D. E. – Fernández, J. R., et al.: Chronotherapy improves blood pressure control and reverts the nondipper pattern in patients with resistant hypertension. *Hypertension*, 2008, 51, s. 69–76, doi:10.1161/HYPERTENSIONAHA.107.096933.
- 15 Hermida, R. C. – Calvo, C. – Ayala, D. E., et al.: Administration-time-dependent effects of doxazosin GITS on ambulatory blood pressure of hypertensive subjects. *Chronobiol Int*, 2004, 21, s. 277–296, dostupné z: <http://www.ncbi.nlm.nih.gov/pubmed/15332347>, vyhledáno 4. 6. 2015.
- 16 Ceral, J. – Solar, M.: Doxazosin: safety and efficacy in the treatment of resistant arterial hypertension. *Blood Press*, 2009, 18, s. 74–77, doi:10.1080/08037050902843726.
- 17 Solar, M. – Ceral, J.: Minoxidil in treatment of severe arterial hypertension. In: *XXIII. výroční sjezd České kardiologické společnosti*. Brno, 2015.
- 18 Kidney, D.: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int Suppl*, 2013, 3, s. 4, doi:10.1038/kisup.2012.76.
- 19 Paiva, L.: Overview of resistant hypertension: A glimpse of the cardiologist's current standpoint. *World J Cardiol*, 2012, 4, s. 275, doi:10.4330/wjcv.4.9.275.
- 20 Bhatt, D. L. – Kandzari, D. E. – O'Neill, W. W., et al.: A controlled trial of renal denervation for resistant hypertension. *N Engl J Med*, 2014, 370, s. 1393–1401, doi:10.1056/NEJMoa1402670.
- 21 Scheffers, I. J. M. – Kroon, A., – Schmidli, J., et al.: Novel baroreflex activation therapy in resistant hypertension: Results of a European multi-center feasibility study. *J Am Coll Cardiol*, 2010, 56, s. 1254–1258, doi:10.1016/j.jacc.2010.03.089.
- 22 Bisognano, J. D. – Bakris, G. – Nadim, M. K., et al.: Baroreflex activation therapy lowers blood pressure in patients with resistant hypertension: Results from the double-blind, randomized, placebo-controlled rheos pivotal trial. *J Am Coll Cardiol*, 2011, 58, s. 765–773, doi:10.1016/j.jacc.2011.06.008.
- 23 Oparil, S. – Schmieder, R. E.: New approaches in the treatment of hypertension. *Circ Res*, 2015, 116, s. 1074–1095, doi:10.1161/CIRCRESAHA.116.303603.
- 24 Lobo, M. D. – Sobotka, P. A. – Stanton, A., et al.: Central arteriovenous anastomosis for the treatment of patients with uncontrolled hypertension (the ROX CONTROL HTN study): a randomised controlled trial. *Lancet*, 2015, 385, s. 1634–1641, doi:10.1016/S0140-6736(14)62053-5.

# Aspirin a clopidogrel – klasický základ protidestičkové léčby

prof. MUDr. Jan Vojáček, DrSc., FESC, FACC

I. interní kardiologická klinika LF UK a FN, Hradec Králové

- 1 Halvorsen, S. – Andreotti, F. – ten Berg, J. M., et al.: Aspirin therapy in primary cardiovascular disease prevention: a position paper of the European Society of Cardiology working group on thrombosis. *J Am Coll Cardiol*, 2014, 64, s. 319–327.
- 2 Baigent, C. – Blackwell, L. – Collins, R., et al.: Antithrombotic Trialists' (ATT) Collaboration. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet*, 2009, 373, s. 1849–1860.
- 3 ISIS-2 (Second International Study of Infarct Survival) Collaborative Group: Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet*, 1988, 13, 2, s. 349–360 (autor neuveden).

# Evolokumab v léčbě dyslipidemií se zaměřením na familiární hypercholesterolemii

doc. MUDr. Michal Vrablík, Ph.D.

Centrum preventivní kardiologie, III. interní klinika 1. LF UK a VFN, Praha

- 1 Seidah, N. G.: PCSK9 as a therapeutic target of dyslipidemia. *Expert Opin Ther Targets*, 2009, 13, s. 19e28.
- 2 Ference, B. A. – Yoo, W. – Walsh, I., et al.: Effect of long-term exposure to lower low-density lipoprotein cholesterol beginning early in life on the risk of coronary heart disease: a Mendelian randomization analysis. *J Am Coll Cardiol*, 2012, 60, s. 2631–2639.
- 3 Cohen, J. C. – Boerwinkle, E. – Mosley, T. H. Jr., et al.: Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med*, 2006, 354, 1264e72.
- 4 Cicero, A. F. G. – Colletti, A. – Borghi, C.: Profile of evolocumab and its potential in the treatment of hyperlipidemia. *Drug Design, Development and Therapy*, 2015, 9, s. 3073–3082.
- 5 Dias, C. S. – Shaywitz, A. J. – Wasserman, S. M., et al.: Effects of AMG145 on low-monoclonal antibody to proprotein convertase subtilisin/kexin type 9 as monotherapy in patients with hypercholesterolemia (MENDEL): a randomised, double-blind, placebo-controlled, phase 2 study. *Lancet*, 2012, 380, s. 1995–2006.
- 6 Kohli, P. – Desai, N. R. – Giugliano, R. P., et al.: Design and rationale of the LAPLACE-TIMI 57 trial: a phase II, double-blind, placebo-controlled study of the efficacy and tolerability of a monoclonal antibody inhibitor of PCSK9 in subjects with hypercholesterolemia on background statin therapy. *Clin Cardiol*, 2012, 35, s. 385–391.
- 7 Blom, D. J. – Hala, T. – Bolognese, M., et al.: A 52-week placebo-controlled trial of evolocumab in hyperlipidemia. *N Engl J Med*, 2014, 370, s. 1809–1819.
- 8 Nordestgaard, B. G. – Chapman, M. J. – Ray, K. K., et al.: Lipoprotein (a) as a cardiovascular risk factor: current status. *European Heart Journal*, 2010, 31, s. 2844–2853.
- 9 Raal, F. – Scott, R. – Somaratne, R., et al.: Low-density lipoprotein cholesterol-lowering effects of AMG 145, a monoclonal antibody to proprotein convertase subtilisin/kexin type 9 serine protease in patients with heterozygous familial hypercholesterolemia: the Reduction of LDL-C with PCSK9 Inhibition in Heterozygous Familial Hypercholesterolemia Disorder (RUTHERFORD) randomized trial. *Circulation*, 2012, 126, s. 2408–2417.
- 10 Raal, F. J. – Honarpour, N. – Blom, D. J., et al.: For the TESLA Investigators. Inhibition of PCSK9 with evolocumab in homozygous familial hypercholesterolemia (TESLA Part B): a randomised, double-blind, placebo-controlled trial. *Lancet*, 2014, 385, s. 341–350.
- 11 Bruckert, E. – Bláha, V. – Stein, E., et al.: Trial assessing long-term use of PCSK9 inhibition in patients with genetic LDL disorders (TAUSSIG). Efficacy and safety in patients with familial hypercholesterolemia receiving lipid apheresis. Poster sessions, AHA Scientific Sessions, Lis-topad 2014, Chicago, s. 15–19.
- 12 Sabatine, M. S. – Giugliano, R. P. – Wiviott, S. D., et al.: Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med*, 2015, 372, s. 1500–1509.

# Novinky v antihypertenzní léčbě

MUDr. Ivan Řiháček, Ph.D. | prof. MUDr. Miroslav Souček, CSc.

II. interní klinika MU a FN u svaté Anny, Brno

- 1 Dostupné z: <http://www.uzis.cz/katalog/roceny/zdravotnicka-rocenka-ceske-republiky-1961-az-2013>; vyhledáno 13. 8. 2016.
- 2 Lancia, G. – Tabard, R. – Narkiewicz, K., et al.: ESH/ESC Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal*, 2013, 34, s. 2159–2219, DOI:10.1093/eurheartj/ehf151.
- 3 Filipovský, J. – Widimský, J. Jr. – Ceral, J., et al.: Diagnostické a léčebné postupy arteriální hypertenze – verze 2012. Doporučení České společnosti pro hypertenzi. *Hypertenze a kardiiovaskulární prevence*, 2012, 3, s. 1–16.
- 4 Cífková, R. – Škodová, Z. – Bruthans, J., et al.: Longitudinální trendy kardiiovaskulární mortality a hodnot krevního tlaku, prevalence, diagnostiky, léčby a kompenzace hypertenze u české populace v letech 1985 až 2007/2008. *J Hypertens (české vydání)*, 2011, 2, s. 3–10.
- 5 Steassen, J. A. – Kuznetsova, T. – Acceptor, R., et al.: OASIS-HT: design of a pharmacogenomic dose-finding study. *Pharmacogenomics*, 2005, 6, s. 755–775.
- 6 Parving, H. H. – Brenner, B. M. – McMurray, J. J. V., et al.: Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *NEJM*, 2012, 367, s. 2204–2213.
- 7 McMurray, J. J. V. – Packer, M. – Desai, A. S., et al.: Angiotensin-neprilysin inhibition versus enalapril in heart failure. *NEJM*, 2014, 371, s. 993–1004.
- 8 Rosa, J. – Widimský, P. – Waldauf, P., et al.: Role of adding spironolactone and renal denervation in true resistant hypertension: one-year outcomes of randomized PRAGUE-15 study. *Hypertension*, 2016, 67, s. 397–403.
- 9 Wright, J. T. – Williamson, P. K. – Snyder, J. K., et al.: A randomised trial of intensive versus standard blood-pressure control. The SPRINT Group. *NEJM*, 2015, 373, s. 2103–2116.
- 10 Van Vark, L. – Bertrand, M. – Akkerhuis, K. M., et al.: Angiotensin-converting enzyme inhibitors reduce mortality in hypertension: a meta-analysis of randomized clinical trials of RAAS inhibitors involving 158 998 patients. *Eur Heart J*, 2012, 33, s. 2088–2097.
- 11 Cheng, J. – Zhang, W. – Zhang, X., et al.: Effect of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on all-cause mortality, cardiovascular deaths, and cardiovascular events in patients with diabetes mellitus. *JAMA Intern Med*, 2014, 174, s. 773–785.

# Novinky v akutní léčbě mozkového infarktu

MUDr. Petr Polidar Neurologické oddělení Thomayerovy nemocnice, Praha  
doc. MUDr. Robert Mikulík, Ph.D. I. neurologická klinika LF MU a FN u sv. Anny, Brno

- Krajcer, Z. – Howell, M. H.: Update on endovascular treatment of peripheral vascular disease: New tools, techniques, and indications. *Tex Heart Inst J*, 2000, 27, s. 369–385.
- Thrift, A. G.: Global stroke statistics. *Int J Stroke*, 2014, 9, s. 6–18.
- Koh, J. S. – Lee, S. J. – Ryu, C. W., et al.: Safety and efficacy of mechanical thrombectomy with solitary stent retrieval for acute ischemic stroke: a systematic review. *Neurointervention*, 2012, 7, s. 1–9.
- Hacke, W. – Kaste, M. – Fieschi, C., et al.: Intravenous thrombolysis with recombinant tissue plasmin activator for acute hemispheric stroke – The European Cooperative Acute Stroke Study (ECASS). *JAMA*, 1995, 274, s. 1017–1025.
- Ciccone, A. – Valvassori, L.: Endovascular treatment for acute ischemic stroke. *N Engl J Med*, 2013, 368, s. 2433–2434.
- Riedel, C. H. – Zimmermann, P. – Jensen-Kondering, U., et al.: The importance of size: successful recanalization by intravenous thrombolysis in acute anterior stroke depends on thrombus length. *Stroke*, 2011, 42, s. 1775–1777.
- Hacke, W. – Kaste, M. – Bluhmki, E., et al.: Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med*, 2008, 359, s. 1317–1329.
- Nour, M. – Scalzo, F. – Liebenskind, D. S.: Ischemia-reperfusion injury in stroke. *Interv Neurol*, 2016, 1, s. 185–199.
- Saver, J. L. – Goyal, M. – Bonafe, A., et al.: Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med*, 2015, 172, s. 2285–2295.
- Mazighi, M. – Mesequer, E. – Labreuche, J., et al.: Bridging therapy in acute ischemic stroke: a systematic review and meta-analysis. *Stroke*, 2012, 43, s. 1302–1308.
- Ding, D.: Endovascular mechanical thrombectomy for acute ischemic stroke: a new standard of care. *J Stroke*, 2015, 17, s. 123–126.
- Powers, M. J. – Derdeyn, C. P. – Biller, J., et al.: American Heart Association Stroke Council 2015 AHA/ASA Focused Update of the 2013 Guidelines for the Early Management of Patients With Acute Ischemic Stroke Regarding Endovascular Treatment. A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*, 2015, 46, s. 3020–3035.
- Gutterman L. R.: Mechanical thrombectomy use in acute stroke treatment. *Endovascular Today*, 2013, 6, s. 43–50.
- Volný, O. – Krajina, A. – Bar, M., et al.: Konsenzus a návrh algoritmu léčby – mechanická trombektomie u akutního mozkového infarktu. *Cesk Slov Neurol N*, 2016, 79, s. 100–110.
- Schönenberger, S. – Möhlenbruch, M. – Pfaff, J., et al.: Sedation vs. intubation for endovascular stroke treatment (SIESTA) – a randomized monocentric trial. *Int J Stroke*, 2015, 10, s. 969–978.
- Mazighi, M. – Mesequer, E. – Labreuche, J., et al.: Bridging therapy in acute ischemic stroke: a systematic review and meta-analysis. *Stroke*, 2012, 43, s. 1302–1308.
- Tagaki, T., et al.: Does intravenous t-PA before acute stroke intervention increase intracranial hemorrhage? – subanalysis of Rescue-japan registry. *Stroke*, 2015, 46.
- MR WITNESS: A study of intravenous thrombolysis with alteplase in MRI-selected patients (MR WITNESS). *ClinicalTrials.gov Identifier: NCT01282242*.
- Amiri, H., et al.: European Cooperative Acute Stroke Study-4: Extending the time for thrombolysis in emergency neurological deficits ECASS-4:EXTEND. *Int J Stroke*, 2016, 11, s. 260–267.
- Huang, X., et al.: Alteplase versus tenecteplase for thrombolysis after ischaemic stroke (ATTEST): A phase 2, randomised, open-label, blinded endpoint study. *Lancet Neurol*, 2015, 14, s. 368–376.

# Reziduální angina pectoris po PCI – kazuistika

MUDr. Jaroslav Brotánek Interní oddělení Thomayerovy nemocnice, Praha

- Želízko, M. – Toušek, F. – Skalický, H.: Summary of the 2013 ESC guidelines on the management of stable coronary artery disease. Prepared by the Czech Society of Cardiology. *Cor et Vasa*, 2014, 56, s. e259–e273.
- Nesukay, E. G.: Treatment of stable angina in Ukraine: CLASSICA study. *Uk J Card*, 2014, 2, s. 43–47.
- Hradec, J.: Co by měl praktický lékař vědět o trimetazidinu? *Kap Kardiol*, 2012, 4, s. 82–88.
- Hradec, J.: Trimetazidin z pohledu klinické praxe. *Farmakoterapie*, 2015, 11, s. 430–434.
- Fragasso, G., et al.: Effect on metabolic modulation by trimetazidine on left ventricular function and phosphocreatine/adenosine triphosphate ratio in patients with heart failure. *Eur Heart J*, 2006, 27, s. 942–948.
- Szwed, H. – Sadowski, Z. – Elikowski, W., et al.: Combination treatment in stable effort angina using trimetazidine and metoprolol: results of a randomised, double-blind, multicentre study (TRIMPOL II). *TRIMetazidine in Poland. Eur Heart J*, 2001, 22, s. 2267–2274.
- Sellier, P. – Brostet, J. P.: Assessment of anti-ischemic and antianginal effect at trough plasma concentration and safety of trimetazidine MR 35 mg in patients with stable angina pectoris: a multicentre, double-blind, placebo-controlled study. *Am J Cardiovasc Drugs*, 2003, 3, s. 361–369.
- Danchin, N. – Marzilli, M. – Parkhomenko, A., et al.: Efficacy comparison of trimetazidine with therapeutic alternatives in stable angina pectoris: a network meta-analysis. *Cardiology*, 2011, 120, s. 59–72.
- Xu, X. – Zhou, Y., et al.: Effect of trimetazidine on recurrent angina pectoris and left ventricular structure in elderly multivessel coronary heart disease patients with diabetes mellitus after drug-eluting stent implantation: A single-centre, prospective, randomized, double-blind study at 2-year follow-up. *Clin Drug Investig*, 2014, 34, s. 251–258.
- Chen, J. – Zhou, S. – Jin, J., et al.: Chronic treatment with trimetazidine after discharge reduces the incidence of restenosis in patients who received coronary stent implantation: A 1-year prospective follow-up study. *Int J Cardiol*, 2014, 174, s. 634–639.
- Gao, D. – Ning, N. – Niu, X., et al.: Trimetazidine: a meta-analysis of randomised controlled trials in heart failure. *Heart*, 2011, 97, s. 278–286.
- Grajek, S. – Michalak, M.: The effect of trimetazidine added to pharmacological treatment on all-cause mortality in patients with systolic heart failure. *Cardiology*, 2015, 131, s. 22–29.
- 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *EHJ*, doi:10.1093/eurheartj/ehw128.
- Yakan, A. S. – Nenaber, C. A.: Pharmacological management of cardiac syndrome Y: A focused review. *Angiol Open Access*, 2015, 3, s. 143, doi: 10.4172/2329-9495.1000143.
- Wang, X. – Nie, S-P.: The coronary slow flow phenomenon: characteristics, mechanisms and implications. *Cardiovasc Diagn Ther*, 2011, 1, s. 37–43.
- Topal, E. – Ozdemir, R. – Barutcu, I., et al.: The effects of trimetazidine on heart rate variability in patients with slow coronary artery flow. *J Electrocardiol*, 2006, 39, s. 211–218.

# Inhibitory ACE versus sartany v léčbě primární hypertenze

MUDr. Jiří Slíva, Ph.D. Ústav farmakologie 3. LF UK, Praha

- Williams, B. – Lacourciere, Y. – Schumacher, H., et al.: Antihypertensive efficacy of telmisartan vs. ramipril over the 24-h dosing period, including the critical early morning hours: a pooled analysis of the PRISMA I and II randomized trials. *J Hum Hypertens*, 2009, 23, s. 610–619.
- Lancia, G. – Tabard, R. – Narkiewicz, K., et al.: ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens*, 2013, 31, s. 1281–1357.
- Li, E. C. – Heran, B. S. – Wright, J. M.: Angiotensin converting enzyme (ACE) inhibitors versus angiotensin receptor blockers for primary hypertension. *Cochrane Database Syst Rev*, 2014, CD009096.
- Bremner, A. D. – Baur, M. – Oddou-Stock, P., et al.: Long-term efficacy and tolerability compared to lisinopril in elderly patients with essential hypertension. *Clin Exper Hypertension*, 1997, 19, s. 1263–1285.
- Barnett, A. H. – Bain, S. C. – Bouter, P., et al.: Angiotensin-receptor blockade versus converting-enzyme inhibition in Type 2 diabetes and nephropathy. *New Eng J Med*, 2004, 351, s. 1952–1961.
- Yusuf, S. – Teo, K. K. – Pogue, J., et al.: Telmisartan, ramipril, or both in patients at high risk for vascular events. *New Eng J Med*, 2008, 358, s. 1547–1559.
- Lacourciere, Y. – Belanger, A. – Godin, C., et al.: Long-term comparison of losartan and enalapril on kidney function in hypertensive type 2 diabetics with early nephropathy. *Kidney International*, 2000, 58, s. 762–769.
- Sozen, A. B. – Kayacan, M. S. – Tansel, T., et al.: Drugs with blocking effects on the renin-angiotensin-aldosterone system do not improve endothelial dysfunction long-term in hypertensive patients. *J Int Med Res*, 2009, 37, s. 996–1002.
- Fogari, R. – Derosa, G. – Ferrari, I., et al.: Effect of valsartan and ramipril on atrial fibrillation recurrence and P-wave dispersion in hypertensive patients with recurrent symptomatic lone atrial fibrillation. *Am J Hypertens*, 2008, 21, s. 1034–1039.
- Fogari, R. – Mugellini, A. – Zoppi, A., et al.: Effect of telmisartan and ramipril on atrial fibrillation recurrence and severity in hypertensive patients with metabolic syndrome and recurrent symptomatic paroxysmal and persistent atrial fibrillation. *J Cardiovasc Pharmacol Ther*, 2011, 17, s. 34–43.
- Fogari, R. – Zoppi, A. – Mugellini, A., et al.: Effects of valsartan or ramipril addition to amlodipine/hydrochlorothiazide combination on left ventricular mass in diabetic hypertensive patients with left ventricular hypertrophy. *Expert Op Pharm*, 2012, 13, s. 1091–1099.
- Spoelstra-de Man, A. M. – van Ittersum, F. J. – Schram, M. T., et al.: Aggressive antihypertensive strategies based on hydrochlorothiazide, candesartan or lisinopril decrease left ventricular mass and improve arterial compliance in patients with type II diabetes mellitus and hypertension. *J Hum Hypertens*, 2006, 20, s. 599–611.
- Bangalore, S. – Fakheri, R. – Toklu, B., et al.: Angiotensin-converting enzyme inhibitors or angiotensin receptor blockers in patients without heart failure? Insights from 254,301 patients from randomized trials. *Mayo Clin Proc*, 2016, 91, s. 51–60.
- Corrao, G. – Zambon, A. – Parodi, A., et al.: Discontinuation of and changes in drug therapy for hypertension among newly-treated patients: a population-based study in Italy. *J Hypertens*, 2008, 26, s. 819–824.

# Chronická pankreatitida – stálá pankreatologická výzva

prof. MUDr. Petr Dítě, DrSc. | MUDr. Martina Bojková Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě

MUDr. Bohuslav Kianička, Ph.D. | MUDr. Tomáš Kupka II. interní klinika FN u sv. Anny, LF MU, Brno

MUDr. Kateřina Kapounková, Ph.D. Katedra podpory zdraví, Fakulta sportovních studií MU, Brno

doc. MUDr. Jana Dvořáčková, Ph.D. Ústav patologie FN, Ostrava

Mgr. Jarmila Šimová | doc. MUDr. Arnošt Martínek, CSc.

CGB laboratoř, Výzkumný institut AGEL, Ostrava

prof. MUDr. Miroslav Souček, CSc. II. interní klinika FN u sv. Anny, LF MU, Brno

- 1 *Chronic pancreatitis – Ha Pan European Pancreatic Club. Guidelines* (v tisku).
- 2 **Yadav, D. – Lowenfels, D.**: The epidemiology of chronic pancreatitis and pancreatic cancer. *Gastroenterology*, 2013, 144, s. 1252–1261.
- 3 **Dítě, P. – Stary, K. – Novotný, I., et al.**: Incidence of chronic pancreatitis in the Czech Republic. *Eur J Gastroent*, 2001, 13, s. 749–750.
- 4 **Etamad, D. – Whitcomb, D.**: Chronic pancreatitis – diagnosis, classification and new genetic development. *Gastroenterology*, 2001, 120, s. 682–707.
- 5 **Whitcomb, D. – Frulloni, L. – Gard, P., et al.**: Chronic pancreatitis. An international draft consensus proposal for a new mechanistic definition. *Pancreatol*, 2016, 16, s. 218–224.
- 6 **Schneider, A. – Lohr, J. M. – Winter, M. W.**: The M-ANHEHEIM classification for chronic pancreatitis: introduction for unifying classification system based on a review of previous classification of the disease. *J Gastroenterol*, 2007, 42, s. 101–119.
- 7 **Perez-Johnston, R. – Saunamäki, N. I. – Sahani, D. V., et al.**: Imaging of chronic pancreatitis (including groove and autoimmune pancreatitis). *Radiol Clin N Amer*, 2012, 50, s. 447–466.
- 8 **Whitcomb, D.**: Genetic risk factors for pancreatic disorders. *Gastroenterology*, 2013, 44, s. 1292–1302.
- 9 **Vlachou, P. A. – Khalil, K. – Jang, H. J., et al.**: IgG<sub>4</sub> related disease: autoimmune pancreatitis and extrapancreatic manifestations. *Radiographics*, 2011, 31, s. 1379–1402.
- 10 **Shimosegawa, T. – Chari, S. T., et al.**: International consensus – diagnostic criteria for autoimmune pancreatitis: guidelines of the International Association of Pancreatology. *Pancreatol*, 2011, 40, s. 352–358.
- 11 **Catalano, M. F. – Sahai, A. – Levy, M., et al.**: EUA-based criteria for the diagnosis of chronic pancreatitis – the Rosemont classification. *Gastrointest Endoscopy*, 2009, 69, s. 1251–1261.
- 12 **Dominguez-Munoz, J. E. – Iglesias-Garcia, J. – Vilarino-Insua, M., et al.**: 13C-mixed triglyceride breath test to assess oral enzyme substitution therapy in patients with chronic pancreatitis. *Clin Gastroenterol Hepatol*, 2007, 5, s. 484–488.
- 13 **Rosendahl, J. – Landt, O. – Bernardova, J., et al.**: CFTR, SPINK1, CTRC, and PRSS1 variants in chronic pancreatitis – is the role of mutated CFTR overestimated? *Gut*, 2013, 62, s. 482–492.
- 14 **Strum, W. B.**: Abstinence of alcohol in chronic pancreatitis. Effect on pain outcome. *J Clin Gastroent*, 1995, 20, s. 37–41.
- 15 **Olesen, S. S.**: Pharmacological pain management in chronic pancreatitis. *WJG*, 2013, 19, s. 7282–7291.
- 16 **Olesen, S. S. – Bouwense, S. A. – Wildr-Smith, O. H., et al.**: Pregabalin reduces pain in patients with chronic pancreatitis in a randomized, controlled trial. *Gastroenterology*, 2011, 141, s. 536–543.
- 17 **Dumoncaeu, J. M. – Delhaye, M. – Tringali, A., et al.**: Endoscopic treatment of chronic pancreatitis: European Society of Gastrointestinal Endoscopy clinical guidelines. *Endoscopy*, 2012, 44, s. 784–800.
- 18 **Winstea, N. S. – Wikcox, C. M.**: Clinical trials of pancreatic enzyme replacement for painful chronic pancreatitis – a review. *Pancreatol*, 2009, 9, s. 344–350.
- 19 **Cahen, G. L. – Gouma, D. J. – Laramee, P., et al.**: Long-term outcomes of endoscopic vs surgical drainage of the pancreatic duct in patients with chronic pancreatitis. *Gastroenterology*, 2011, 141, s. 1690–1695.
- 20 **Dítě, P. – Růžicka, M. – Zboril, V., et al.**: A prospective, randomized trial comparing endoscopic and surgical therapy for chronic pancreatitis. *Endoscopy*, 2003, 35, s. 553–558.
- 21 **Silkens, E. C. M. – Cahen, D. – Kuipers, E. J., et al.**: Pancreatic enzyme replacement therapy in chronic pancreatitis. *Best Clin Practise Clin Gastroenterol*, 2010, s. 337–347.
- 22 **Berry, A. M.**: Pancreatic enzyme replacement therapy during pancreatic insufficiency. *Nutr In Clin Practice*, 2014, 29, s. 312–321.
- 23 **Meyer, J. H. – Elhoff, J. – Porter-Fink, V., et al.**: Human postprandial gastric emptying of 1–3 millimeter spheres. *Gastroenterology*, 1988, 94, s. 1215–1225.
- 24 **Lindkvist, B.**: Diagnosis and treatment of pancreatic exocrine insufficiency. *WJG*, 2013, 19, s. 7258–7266.
- 25 **Trang, T. – Chan, J. – Graham, D. Y.**: Pancreatic enzyme replacement therapy for pancreatic exocrine insufficiency in the 21st century. *WJG*, 2014, 7, s. 11467–11483.
- 26 **Sickens, E. C. – Cahen, D. L. – Koch, A. D., et al.**: The prevalence of fat-soluble vitamins deficiencies and decrease of bone mass in patients with chronic pancreatitis. *Pancreatol*, 2013, 13, s. 238–242.
- 27 **Keller, J. – Buckem, S. – Jahr, C., et al.**: A modified 13-C breath test detected moderate pancreatic insufficiency. *Pancreas*, 2011, 40, s. 1201–1205.

## Pokroky v diagnostice a terapii akutní pankreatitidy

prof. MUDr. Petr Dítě, DrSc. | MUDr. Martina Bojková

Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě

Jarmila Šimová CGB laboratoř, a. s., Ostrava, člen skupiny Agel

MUDr. Bohuslav Kianička, Ph. D. II. interní klinika FN u sv. Anny, LF MU, Brno

MUDr. Tomáš Kupka | Mgr. Lenka Dovrtělová, Ph.D. | doc. MUDr. Arnošt Martínek, CSc.

Interní klinika – gastroenterologické oddělení FN Ostrava, LF Ostravské univerzity v Ostravě

prof. MUDr. Miroslav Souček, CSc. II. interní klinika FN u sv. Anny, LF MU, Brno

- 1 **Špičák, J.**: *Akutní pankreatitida*. Grada, Praha, 2005.
- 2 **Criddle, D. N.**: The role of fat and alcohol in acute pancreatitis: a dangerous lesion. *Pancreatol*, 2015, 15 (dopl. 4), s. S6–S12.
- 3 **Dumoncaeu, J. M. – Andriulli, A. – Elmunzer, J. B., et al.**: Prophylaxis of post – ERCP pancreatitis: ESGE guideline. *Endoscopy*, 2014, 46, s. 799–815.
- 4 **Tender, S.**: Drug induced acute pancreatitis-does it exist? *WJG*, 2014, 20, s. 16529–16534.
- 5 **Vinklerová, J. – Grygar, J. – Procházka, V., et al.**: Je indukovaná akutní pankreatitida retrospektivní analýza. *Česká a Slo Gastroenterol*, 2008, 62 (dopl. 3), s. 31–35.
- 6 **Whitcomb, D. C.**: Clinical practice—Acute pancreatitis. *N Engl J Med*, 2006, 354, s. 2142–2150.
- 7 **Arvanitakis, M. – Delhaye, M. – DeMaertelaere, J., et al.**: Computed tomography and magnetic resonance imaging in the assessment of acute pancreatitis. *Gastroenterology*, 2004, 126, s. 715–723.
- 8 **Bollen, T.**: Imaging of acute pancreatitis: update of revised Atlanta classification *RADIOL. Clin N Amer*, 2012, 50, s. 429–450.
- 9 **Baron, T. H.**: Imaging of severe acute pancreatitis. *Cleve Clin J Med*, 2013, 80, s. 354–359.
- 10 **Baltazar, E. J.**: Acute pancreatitis: assesment of severity with clinical and CT evaluation. *Radiology*, 2002, 223, s. 603–613.
- 11 **Banks, P. A. – Bollen, T. L. – Dervenis, C., et al.**: Classification of acute pancreatitis 2012: revision of the Atlanta classification and definitions by international consensus. *Gut*, 2013, 62, s. 102–111.
- 12 **Brown, A. – Orav, J. – Banks, A.**: Hemoconcentration is a nearly marker for organ failure and necrotizing pancreatitis. *Pancreas*, 2000, 20, s. 367–370.
- 13 **Zerem, E.**: Treatment of severe acute pancreatitis and its complications. *WJG*, 2014, 14, s. 1879–1899.
- 14 **Basurto, O. X. – Rigau, C. E. – Urutia, G., et al.**: Opioids for acute pancreatitis pain. *Cochrane Database Syst Rev*, 2013, 26, 7, CD 009179.
- 15 **De Vries, A. C. – Besselink, M. G. – Buskens, E., et al.**: Randomized controlled trial of antibiotic prophylaxis in severe acute pancreatitis: relationship between methodological quality and outcome. *Pancreatol*, 2007, 75, s. 531–538.
- 16 **Hart, P. A. – Bechtold, M. L. – Marshall, J. B., et al.**: Prophylactic antibiotics in necrotizing pancreatitis. *South Med J*, 2008, 101, s. 1126–1131.
- 17 **Gianotti, L. – Meier, R. – Lobo, D. N., et al.**: ESPEN Guidelines on Parenteral Nutrition: Pancreas. *Clin Nutr*, 2009, 28, s. 428–435.
- 18 **Al-Omran, M. – Albalawi, Z. H. – Tashkandi, M. F., et al.**: Enteral versus parenteral nutrition for acute pancreatitis. *Cochrane Database Syst Rev*, 20. 1. 2010, s. 1.
- 19 **Andrašina, T. – Bohatá, Š.**: Zobrazovací metody u akutní pankreatitidy. *Klinická pankreatologie*, 2016, s. 130–138.
- 20 **Šenkyřík, M.**: Akutní pankreatitida – základy konzervativní léčby. *Klinická pankreatologie*, 2016, s. 55–80.

# Diagnostika a příznaky Fabryho choroby

MUDr. Lubor Golář II. interní klinika kardiologie a angiologie 1. LF UK a VFN, Praha

- Spada, M. – Pagliardini, S. – Yasuda, M., et al.: High incidence of later-onset Fabry disease revealed by newborn screening. *Am J Hum Genet*, 2006, 79, s. 31–40.
- Mechtler, T. P. – Stary, S. – Metz, T. F., et al.: Neonatal screening for lysosomal storage disorders: feasibility and incidence from a nationwide study in Austria. *Lancet*, 2012, 379, s. 335–341.
- Lin, H. Y. – Chiny, K. W. – Hus, J. H. – Yu, H. C., et al.: High incidence of the cardiac variant of Fabry disease revealed by newborn screening in the Taiwan Chinese population. *Circ Cardiovasc Genet*, 2009, 2, s. 450–456.
- Paleček, T. – Honzík, J. – Poupátová, H., et al.: Prevalence of Fabry disease in male patients with unexplained left ventricular hypertrophy in primary cardiology practice: prospective Fabry cardiomyopathy screening study (FACSS). *J Inher Metab Dis*, 2014, 37, s. 455–460.
- Elliot, P. – Baker, R. – Pasquale, F., et al.: ACES study group. Prevalence of Anderson-Fabry disease in patients with hypertrophic cardiomyopathy: the European Anderson-Fabry Disease survey. *Heart*, 2011, 97, s. 1957–1960.
- Merta, M. – Reiterová, J. – Ledvinová, J., et al.: A nationwide blood spot screening study for Fabry disease in the Czech Republic haemodialysis patient population. *Nephrol Dial Transplant*, 2007, 22, s. 179–186.
- Linhart, A. – Paleček, T. – Bultas, J., et al.: New insights in cardiac structural changes in patients with Fabry's disease. *Am Heart J*, 2000, 139, 6, s. 1101–1108.
- Shah, J. S. – Hughes, D. A. – Sachdev, B., et al.: Prevalence and clinical significance of cardiac arrhythmia in Anderson-Fabry disease. *Am J Cardiol*, 2005, 96, s. 842–846.
- Alroy, J. – Sharda, S. – Kopp, J. B.: Renal pathology in Fabry disease. *J Am Soc Nephrol*, 2002, 13, s. S134–S138.
- Rofls, A. – Bottche, T. – Zschiesche, M., et al.: Prevalence of Fabry disease in patients with cryptogenic stroke: a prospective study. *Lancet*, 2005, 366, s. 1794–1796.
- Chamoles, N. – Blanco, M. – Gaglioli, D.: Fabry disease: enzymatic diagnosis in dried blood spots on filter paper. *Clin Chim Acta*, 2001, 308, s. 195–196.
- Thurberg, B. L. – Politei, J. M.: Histologic abnormalities of placental tissues in Fabry disease: a case report and review of the literature. *Hum Pathol*, 2012, 43, s. 610–614.
- Eng, C. M. – Guffon, N. – Wilcox, W. R., et al.: Safety and efficacy of recombinant human alpha-galactosidase A – replacement therapy in Fabry's disease. *N Engl J Med*, 2001, 345, s. 9–16.
- Banikazemi, M. – Bultas, J. – Waldek, S., et al.: Agalsidase-beta therapy for advanced Fabry disease: a randomized trial. *Ann Intern Med*, 2007, 146, s. 77–86.
- Schiffmann, R. – Kopp, J. B. – Austin, 3<sup>rd</sup> H. A., et al.: Enzyme replacement therapy in Fabry disease: a randomized controlled trial. *JAMA*, 2001, 285, s. 2743–2749.
- Kampmann, C. – Linhart, A. – Devereux, R. B., et al.: Effect of agalsidase alfa replacement therapy on Fabry disease-related hypertrophic cardiomyopathy: a 12- to 36-month, retrospective, blinded echocardiographic pooled analysis. *Clin Ther*, 2009, 31, s. 1966–1976.
- Ojo, A. – Meier-Kriesche, H. U. – Friedman, G., et al.: Excellent outcome of renal transplantation in patients with Fabry's disease. *Transplantation*, 2000, 69, s. 2337–2339.

## Léčba CHOPN – současnost a novinky

doc. MUDr. Norbert Pauk, Ph.D.

Klinika pneumologie a hrudní chirurgie Nemocnice Na Bulovce a 3. LF UK, Praha

- Global strategy for diagnosis, management and prevention of chronic pulmonary disease. *GOLD Report*, revidováno 2016, dostupné z: [www.goldcopd.org](http://www.goldcopd.org).
- Woodruff, P. G. – Agusti, A. – Roche, N., et al.: Current concepts in targeting chronic obstructive pulmonary disease pharmacotherapy: making progress towards personalised management. *Lancet*, 2015, 385, s. 1789–1798.
- Koblížek, V. – Chlumský, J. – Zindr, V., et al.: *Doporučený postup ČPFS pro diagnostiku a léčbu stabilní CHOPN*. Praha, Maxdorf Jessenius, 2013, s. 134.
- Kašák, V.: Inhalční systémy v terapii astmatu a chronické obstrukční plicní nemoci. *Remedia*, 2014, 24, s. 315–320.
- Diagnosis of diseases of chronic airflow limitation: Asthma, COPD and Asthma-COPD Overlap Syndrome (ACOS)*. GINA 2014, GOLD 2014, dostupné z: [www.ginasthma.org](http://www.ginasthma.org).
- Jadwiga, A. – Wedzicha, M. D. – Donald Banerji, M. D., et al.: Indacaterol-glycopyrronium versus salmeterol-fluticasone for COPD. *NEJM*, 15. 5. 2016, DOI: 10.1056/NEJMo1516385.

## Léčba astmatu – současnost a novinky

doc. MUDr. Norbert Pauk, Ph.D.

Klinika pneumologie a hrudní chirurgie Nemocnice Na Bulovce a 3. LF UK, Praha

- GINA (Global Initiative for Asthma). *Global Strategy for Asthma Management and Prevention*, 2015.
- ČIPA (Česká iniciativa pro astma): *Strategie diagnostiky, prevence a léčby astmatu. Uvedení globální strategie do praxe v ČR*. Praha, Jalna, 2012.
- Teřil, M.: Diagnostický přístup k astmatu prizmatem eozinofilie a alergie. *Stud Pneumol Phthiseol*, 2009, 69, s. 130–139.
- Kašák, V.: *Asthma bronchiale*. Praha, Maxdorf, 2013.
- Kašák, V. – Feketeová, E.: Vliv nesprávné inhalační techniky na úroveň kontroly nad astmatem. *Alergie*, 2010, 12, s. 244–257.
- Teřil, M. – Čáp, P. – Dvořáková, R., et al.: *Doporučený postup diagnostiky a léčby bronchiálního astmatu*. Semily, GEUM, 2015.
- Teřil, M.: Léčba astmatu – chyby a omyly každodenní praxe. *Med Praxi*, 2013, 10, s. 97–103.
- Sedlák, V. – Chlumský, J. – Teřil, M., et al.: *Doporučený postup diagnostiky a léčby obtížně léčitelného bronchiálního astmatu*. 2011, dostupné z: <http://www.pneumologie.cz/guidelines/>, vyhledáno 30. 5. 2016.

## Novinky v léčbě metabolického syndromu

prof. MUDr. Štěpán Svačina, DrSc., MBA III. interní klinika 1. LF UK a VFN, Praha

- Svačina, Š.: *Obezitologie a teorie metabolického syndromu*. Triton, Praha, 2013.
- Ferrannini, E. – Muscilli, E. – Frascerra, S., et al.: Metabolic response to sodium-glucose cotransporter 2 inhibition in type 2 diabetic patients. *J Clin Invest*, 2014, 124, s. 499–508.
- Nissen, S. E. – Wolski, K. E. – Prcela, L., et al.: Effect of naltrexone-bupropion on major adverse cardiovascular events in overweight and obese patients with cardiovascular risk factors: a randomized clinical trial. *JAMA*, 2016, 8, 315, s. 990–1004.
- Feinstein, M. J. – Lloyd-Jones, D. M.: Monoclonal antibodies for lipid management. *Curr Atheroscler Rep*, 2016, 18, s. 39, e-publikace.
- Ferri, N. – Ruscica, M.: Proprotein convertase subtilisin/kexin type 9 (PCSK9) and metabolic syndrome: insights on insulin resistance, inflammation, and atherogenic dyslipidemia. *Endocrine*, 1. 4. 2016, e-publikace.
- Zinman, B. – Wanner, Ch. – Lachinet, J. M., et al.: EMPA-REG OUTCOME investigators: Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med*, 2015, 373, s. 2117–2128.
- Wanner, C. – Inzucchi, S. E. – Lachin, J. M., et al.: EMPA-REG OUTCOME investigators: Empagliflozin and progression of kidney disease in type 2 diabetes. *N Engl J Med*, 2016, 375, s. 323–334.
- Marso, S. P. – Daniels, G. H. – Brown-Frandsen, K., et al.: LEADER Trial investigators: Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med*, 2016, 375, s. 311–322.



# Fixní kombinace bazálního inzulínového analoga a GLP-1 po selhání bazálního inzulínu

prof. MUDr. Zdeněk Rušavý, Ph.D. | MUDr. Michal Žourek I. interní klinika FN a LF UK, Plzeň

- 1 Barnett, A. H.: Tolerability and efficacy exenatide and titrated insulin glargine in adult patients with type 2 diabetes previously uncontrolled with metformin or a sulfonylurea: a multinational, randomised, open-label, two-period, crossover, noninferiority trial. *Clin Ther*, 2007, 29, s. 2333–2348.
- 2 Hein, G. J. – Baker, Ch. – Hsieh, J., et al.: GLP-1 and GLP-2 – stimulated postprandial lipemia in normal and insulin-resistant states. *Diabetes*, 2013, 62, s. 373–381.
- 3 Jeppesen, P. B. – Gildou, R. – Pertkiewicz, M., et al.: Randomised placebo-controlled trial of teduglutide in reducing parenteral nutrition and/or intravenous fluid requirements in patients with short bowel syndrome. *Gut*, 2011, 60, s. 902–914.
- 4 Davies, M. – Bain, S. – Atkin, S., et al.: Efficacy and safety of liraglutide versus placebo as add-on to glucose-lowering therapy in patients with type 2 diabetes and moderate renal impairment (LIRA-RENAL): A randomized clinical trial. *Diabetes Care*, 2016, 39, s. 222–230.
- 5 Retnakaran, R. – Kramer, C. – Zinman, B., et al.: Liraglutide and the preservation of pancreatic  $\beta$ -cell function in early type 2 diabetes. The LIBRA trial. *Diabetes Care*, 2014, 37, s. 3270–3278.
- 6 Marso, S. – Daniels, G. – Zinman, B., et al.: Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med*, 2016, 375, s. 311–322.
- 7 Eng, C. – Kramer, C. – Zinman, B. – Retnakaran, R.: Glucagon-like peptide-1 receptor agonist and basal insulin combination treatment for the management of type 2 diabetes: a systematic review and meta-analysis. *Lancet*, 2014, 384, s. 2228–2234.
- 8 Cramer, J. – Pugh, M. J.: The influence of insulin use on glycemic control. How well do adults follow prescriptions for insulin? *Diabetes Care*, 2005, 28, s. 78–83.
- 9 Svanum, H. A. – Lage, M. J., et al.: Early discontinuation and restart of insulin in the treatment of type 2 diabetes mellitus. *Diabetes Ther*, 2014, 5, s. 225–242.
- 10 Gough, S. – Bode, B. – Buse, J., et al.: Efficacy and safety of a fixed-ratio combination of insulin degludec and liraglutide (IDegLira) compared with its components given alone results of a phase 3, open label, randomised, 26 week treat-to-target trial in insulin naïve patients with type 2 diabetes. *Lancet Diabetes Endocrinol*, 2014, 2, s. 885–893.
- 11 Buse, J. – Vilsbøll, T. – Thurman, J., et al.: Contribution of liraglutide in the fixed-ratio combination of insulin degludec and liraglutide (IDegLira): *Diabetes Care*, 2014, 37, s. 2926–2933.
- 12 Lingvay, I. – Manghi, F. – García-Hernández, P., et al.: Effect of insulin glargine up-titration vs insulin degludec/liraglutide on glycated hemoglobin levels in patients with uncontrolled type 2 diabetes. The dual v randomized clinical trial. *JAMA*, 2016, 315, s. 898 – 907.
- 13 The FLAT-Sugar trial investigators: Glucose variability in a 26-week randomized comparison of mealtime treatment with rapid-acting insulin versus GLP-1 agonist in participants with type 2 diabetes at high cardiovascular risk. *Diab Care*, 2016, 39, s. 973–981.
- 14 Bajaj, H. – Venn, K. – Ye, Ch., et al.: Least glucose variability is observed with the combination of a GLP-1 receptor agonist and basal insulin among four commonly used insulin regimen in type 2 diabetes (VARIATION Study). *Canadian J Diabet*, 2014, dopl.: s. S41.

## Novinky v léčbě diabetika 2. typu

prof. MUDr. Jan Škrha, DrSc., MBA

III. interní klinika – klinika endokrinologie a metabolismu, 1. LF UK a VFN, Praha

- 1 Škrha, J.: Současné postavení inhibitorů SGLT2 v léčbě diabetu 2. typu. *Remedia*, 2016, 26, s. 1–5.
- 2 Zinman, B. – Wanner, Ch. – Lachin, J. M., et al.: Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med*, 2015, 373, s. 2117–2128.
- 3 Uccellatore, A. – Genovese, S. – Dicembrini, I., et al.: Comparison review of short-acting and long-acting glucagon-like peptide-1 receptor agonists. *Diabetes Ther*, 2015, 6, s. 239–256.
- 4 Guo, X. H.: The value of short- and long-acting glucagon-like peptide-1 agonists in the management of type 2 diabetes mellitus: experience with exenatide. *Curr Med Res Opin*, 2016, 32, s. 61–76.
- 5 Green, J. B. – Bethel, M. A., et al.: Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med*, 2015, 373, s. 232–242.
- 6 White, W. B. – Cannon, C. P. – Heller, S. R., et al.: Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med*, 2013, 369, s. 1327–1335.
- 7 Bentley-Lewis, R. – Aguilar, D. – Riddle, M. C., et al.: Rationale, design, and baseline characteristics in Evaluation of LIXisenatide in Acute Coronary Syndrome, a long-term cardiovascular end point trial of lixisenatide versus placebo. *Am Heart J*, 2015, 169, s. 631–638.
- 8 Scirica, B. M. – Bhatt, D. L. – Braunwald, E., et al.: Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med*, 2013, 369, s. 1317–1326.
- 9 Bonora, E. – Cigolini, M.: DPP-4 inhibitors and cardiovascular disease in type 2 diabetes mellitus. Expectations, observations and perspectives. *Nutr Metab Cardiovasc Dis*, 2016, 26, s. 273–284.
- 10 Deacon, C. F. – Lebovitz, H. E.: Comparative review of dipeptidyl peptidase-4 inhibitors and sulfonylureas. *Diab Obes Metab*, 2016, 18, s. 333–347.
- 11 Buse, J. B. – Rodbard, H. W. – Serrano, C. T., et al.: Randomized clinical trial comparing basal insulin peglispro and insulin glargine in patients with type 2 diabetes previously treated with basal insulin: IMAGINE 5. *Diabetes Care*, 2016, 39, s. 92–100.
- 12 Keller, S. – Mathieu, C. – Kapur, R., et al.: A meta-analysis of rate ratios for nocturnal confirmed hypoglycaemia with insulin degludec vs. insulin glargine using different definitions for hypoglycaemia. *Diabetic Med*, 2016, 33, s. 478–487.
- 13 Home, P. D. – Yki-Jarvinen, H. – Bolli, G. B., et al.: New insulin glargine 300U/ml in type 2 diabetes using basal plus meal-time insulin: sustained glycaemic control and less hypoglycaemia compared with glargine 100U/ml (EDITION 1). *Diabetic Med*, 2015, 32, dopl. 1, s. 78.
- 14 de la Pena, A. – Seger, M. – Soon, D., et al.: Bioequivalence and comparative pharmacodynamics of insulin lispro 200U/mL relative to insulin lispro (Humalog (R)) 100 U/mL. *Clin Pharmacol Drug Develop*, 2016, 5, s. 69–75.
- 15 Lokhandwala, T. – Smith, N. – Sternhufvud, C., et al.: A retrospective study of persistence, adherence, and health economic outcomes of fixed-dose combination vs loose-dose combination of oral anti-diabetes drugs. *J Med Econom*, 2016, 19, s. 203–212.
- 16 Perez, A. – Jacks, R. – Arora, V. – Spanheimer, R.: Effects of pioglitazone and metformin fixed-dose combination therapy on cardiovascular risk markers of inflammation and lipid profile compared with pioglitazone and metformin monotherapy in patients with type 2 diabetes. *J Clin Hypertension*, 2010, 12, s. 973–982.
- 17 Scheen, A.: Linagliptin plus metformin: a pharmacokinetic and pharmacodynamic evaluation. *Exp Opin Drug Metab Toxicol*, 2013, 9, s. 363–377.
- 18 Tan, X. Y. – Hu, J. B.: Combination therapy for type 2 diabetes: dapagliflozin plus metformin. *Exp Opin Pharmacother*, 2016, 17, s. 117–126.
- 19 Aronson, R.: Single-pill combination therapy for type 2 diabetes mellitus: linagliptin plus empagliflozin. *Curr Med Res Opin*, 2015, 31, s. 901–911.
- 20 Morales, J. – Merker, L.: Minimizing hypoglycemia and weight gain with intensive glucose control: Potential benefits of a new combination therapy (IDegLira). *Adv Ther*, 2015, 32, s. 391–403.
- 21 *Doporučený postup péče o diabetes mellitus 2. typu. Doporučení České diabetologické společnosti ČLS JEP.* Dostupné z: [www.diab.cz](http://www.diab.cz)

## DM2T: léčebný „switch“ pomocí dapagliflozinu, snížení hmotnosti jako přidaná hodnota – kazuistika

MUDr. David Bartoš Diabetologická ambulance, Centrum zdraví Sagena, Frýdek-Místek

Diabetologická ambulance, Vítkovická nemocnice, a. s.

- 1 Haluzík, M. – Rychlík, I. – Franěk, M.: *Levdiny a léčba diabetu: současná a perspektivy terapie glifloziny*. Praha, Mladá fronta, 2016.
- 2 Škrha, J., et al.: *Diabetologie*. Praha, Galén, 2009.
- 3 Perušičová, J., et al.: Studie MOTUDI – ovlivnění hmotnosti inzulínovou léčbou u nemocných s DM2T v ČR. *Klinická farmakologie a farmacie*, 2012, 26, s. 50–54.

# Současné možnosti léčby obezity a zdravotně závažné nadváhy

MUDr. Petr Sucharda, CSc. III. interní klinika VFN a 1. LF UK, Praha

- Enzi, G. – Busetto, L. – Inelmen, E. M., et al.: Historical perspective: visceral obesity and related comorbidity in Joannes Baptista Morgagni's 'De sedibus et causis morborum per anatomem indagata'. *Int J Obes Relat Metab Disord*, 2003, 27, s. 534–535.
- World Health Organisation: *Fact sheet No. 311* (aktualizováno v lednu 2015). Dostupné z: [www.who.int/mediacentre/factsheets/fs311/en/](http://www.who.int/mediacentre/factsheets/fs311/en/), vyhledáno 6. 6. 2016.
- Ahima, R. S. – Mitchell, A. L.: The health risk of obesity – better metrics imperative. *Science*, 2013, 341, s. 856.
- Dixon, J. B. – Lambert, G. W.: The obesity paradox a reality that requires explanation and clinical interpretation. *Atherosclerosis*, 2013, 226, s. 47–48.
- Pontzer, H. – Raichlen, D. A. – Wood, B. M., et al.: Hunter-gatherer energetics and human obesity. *PLoS One*, 2012, 7, s. e40503.
- Berbesque, J. C. – Marlowe, F. W. – Shaw, P., et al.: Hunter-gatherers have less famine than agriculturalists. *Biol Lett*, 2010, 20130853, doi: 10.1098/rsbl.2013.0853.
- Semper, H. M. – Povey, R. – Clark-Carter, D.: A systematic review of the effectiveness of smartphone applications that encourage dietary self-regulatory strategies for weight loss in overweight and obese adults. *Obes Rev*, 18. 5. 2016, doi:10.1111/obr.12428, Epub před tiskem.
- [http://www.vstj.cz/?pg=rekonidnci\\_centra](http://www.vstj.cz/?pg=rekonidnci_centra), vyhledáno 16. 6. 2016.
- Hainer, V. – Toplak, H. – Mittraku, A.: Treatment modalities of obesity: what fits whom? *Diabetes Care*, 2008, 31, dopl. 2, s. S269–S277.
- MacDaniels, J. S. – Schwarz, T. L.: Effectiveness, tolerability and practical application of the newer generation anti-obesity medications. *Drugs Context*, 2016, 8, 212291.
- Apovian, C. M. – Aronne, L. – Rubino, L., et al.: A randomized, phase 3 trial of naltrexone SR/ bupropion SR on weight and obesity-related risk factors (COR-II). *Obesity*, 2013, 21, s. 935–943.
- Fidler, M. C. – Sanchez, M. – Raether, B., et al.: BLOSSOM Clinical Trial Group. A one-year randomized trial of lorcaserin for weight loss in obese and overweight adults: the BLOSSOM trial. *J Clin Endocrinol Metab*, 2011, 96, s. 3067–3077.
- Martin, C. K. – Redman, L. M. – Zhang, J., et al.: Lorcaserin, a 5-HT(2C) receptor agonist, reduces body weight by decreasing energy intake without influencing energy expenditure. *J Clin Endocrinol Metab*, 2011, 96, s. 837–845.
- Toplak, H. – Woodward, E. – Yumuk, V., et al.: 2014 EASO Position statement on the use of anti-obesity drugs. *Obes Facts*, 2015, 8, s. 166–174.
- Krentz, A. J. – Fujioka, K. – Hompesch, M.: Evolution of pharmacological obesity treatments: focus on adverse side-effect profiles. *Diabetes Obes Metab*, 2016, 18, s. 558–570.
- Gadde, K. M. – Looping, M. F. – Wagner, H. R. 2<sup>nd</sup>, et al.: Zonisamide for weight reduction in obese adults: a 1-year randomized controlled trial. *Arch Intern Med*, 2012, 172, s. 1557–1564.
- Fujioka, K.: Current and emerging medications for overweight or obesity in people with comorbidities. *Diabetes Obes Metab*, 2015, 17, s. 1021–1032.
- Gauner, R.: Comparative efficiency and safety of pharmacological approaches to the management of obesity. *Diabetes Care*, 2011, 34, dopl. 2, s. S349–S354.
- Georgie, M. – Rajaram, M. – Shanmugam, E.: New and emerging drug molecules against obesity. *J Cardiovasc Pharmacol Ther*, 2014, 19, s. 65–76.
- Joharapurkar, A. A. – Dhanesha, N. A. – Jain, M. R.: Inhibition of the methionine aminopeptidase 2 enzyme for the treatment of obesity. *Diabetes Metab Syndr Obes*, 2014, 7, s. 73–84.
- Kim, D. D. – Krishnarajah, J. – Lillioja, S., et al.: Efficacy and safety of belorabin for weight loss in obese adults: a randomized controlled trial. *Diabetes Obes Metab*, 2015, 17, s. 566–572.
- Kurylowicz, A.: In search of new therapeutic targets in obesity treatment: Sirtuins. *Int J Mol Sci*, 2016, 17, 4.
- Nguyen, L. T. – Chen, H. – Pollock, C. A., et al.: Sirtuins-mediators of maternal obesity-induced complications in offspring? *FASEB J*, 2016, 30, s. 1383–1390.
- Salehi, P. – Hsu, I. – Azen, C. G., et al.: Effects of exenatide on weight and appetite in overweight adolescents and young adults with Prader-Willi syndrome. *Pediatr Obes*, 13. 4. 2016, doi: 10.1111/ajpo.12131, Epub před tiskem.
- Kouno, T. – Akiyama, N. – Ito, T., et al.: Ghrelin O-acyltransferase knockout mice show resistance to obesity when fed high-sucrose diet. *J Endocrinol*, 2016, 228, s. 115–125.
- Miller, T. D. – Nogueiras, R. – Andermann, M. L., et al.: Ghrelin. *Mol Metab*, 2015, 4, s. 437–460.
- Skow, M. A. – Bergmann, N. C. – Knop, F. K.: Diabetes and obesity treatment based on dual incretin receptor activation – "twincretins". *Diabetes Obes Metab*, 10. 5. 2016, doi: 10.1111/dom.12685, Epub před tiskem.
- Doležalová, K. – Býma, S. – Fried, M. – Svačina, Š., et al.: *Bariatrická chirurgie a primární péče*. Axonite, Praha, 2012.
- Fried, M., et al.: *Bariatrická a metabolická chirurgie. Nové postupy v léčbě obezity a metabolických poruch*. Mladá fronta, Praha, 2011.
- Pories, W. J. – Swanson, M. S. – MacDonald, K. G., et al.: Who would have thought it? An operation proves to be the most effective therapy for adult-onset diabetes mellitus. *Ann Surg*, 1995, 222, s. 339–350.
- Sucharda, P.: Hormonální a metabolické důsledky bariatrických výkonů. In: Doležalová, K. – Býma, S. – Fried, M. – Svačina, Š., et al.: *Bariatrická chirurgie a primární péče*. Axonite, Praha, 2012, s. 89–98.
- Fried, M. – Tumák, V. – Kopert, J. M., et al.: a v zastoupení evropské sekce Mezinárodní federace bariatrické a metabolické chirurgie (IFSO-EC) a Evropské obezitologické společnosti (EASO). *Interdisciplinární evropské doporučení metabolické a bariatrické chirurgie*. 2013, český překlad dostupný z: [http://www.obesitas.cz/download/interdisciplinarni\\_evropska\\_doporučení\\_metabolické\\_a\\_bariatrické\\_chirurgie.pdf](http://www.obesitas.cz/download/interdisciplinarni_evropska_doporučení_metabolické_a_bariatrické_chirurgie.pdf), vyhledáno 16. 6. 2016.
- Tumák, V. – Tsigos, C. – Fried, M., et al.: Obesity Management Task Force of the European Association for the Study of Obesity. European guidelines for obesity management in adults. *Obes Facts*, 2015, 8, s. 402–424.
- Rubino, F. – Nathan, D. M. – Eckel, R. H., et al.: Delegates of the 2<sup>nd</sup> Diabetes Surgery Summit. Metabolic surgery in the treatment algorithm for type 2 diabetes: A joint statement by international diabetes organizations. *Diabetes Care*, 2016, 39, s. 861–877.
- Cummings, D. E. – Cohen, R. V.: Bariatric/metabolic surgery to treat type 2 diabetes in patients with a BMI < 35 kg/m<sup>2</sup>. *Diabetes Care*, 2016, 39, s. 924–933.
- Lupoli, R. – Di Minno, M. N. – Guidone, C., et al.: Effects of bariatric surgery on markers of subclinical atherosclerosis and endothelial function: a meta-analysis of literature studies. *Int J Obes (Lond)*, 2016, 40, s. 395–402.
- Rohde, U. – Hedbäck, N. – Fluid, L. L., et al.: Effect of the EndoBarrier Gastrointestinal Liner on obesity and type 2 diabetes: a systematic review and meta-analysis. *Diabetes Obes Metab*, 2016, 18, s. 300–305.
- Gaur, S. – Levy, S. – Mathus-Vliegen, L., et al.: Balancing risk and reward: a critical review of the intragastric balloon for weight loss. *Gastrointest Endosc*, 2015, 81, s. 1330–1336.
- Sanmiguel, C. P. – Conklin, J. L. – Cunneen, S. A., et al.: Gastric electrical stimulation with the TANTALUS System in obese type 2 diabetes patients: effect on weight and glycaemic control. *J Diabetes Sci Technol*, 2009, 3, s. 964–970.

## Časná léčba inzulinem u pacientů s diabetes mellitus 2. typu

MUDr. Milan Flekač, Ph.D. | MUDr. Jan Šoupal III. interní klinika VFN a 1. LF UK, Praha

- Lindstrom, J. – Ilanne-Parikka, P. – Peptonen, M., et al.: Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet*, 2006, 368, s. 1673–1679.
- Knowler, W. C. – Fowler, S. E. – Hamman, R. F., et al.: 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*, 2009, 374, s. 1677–1686.
- The ORIGIN Trial Investigators: Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med*, 2012, 367, s. 319–328, DOI: 10.1056/NEJMoa1203858.
- Kahn, S. E. – Haffner, S. M. – Heise, M. A., et al.: Glycemic durability of rosiglitazone, metformin, or glyburide monotherapy. *N Engl J Med*, 2006, 355, s. 2427–2443.
- American Diabetes Association: Approaches to glycaemic treatment. *Diabetes Care*, 2015, 38, s. S41–S48.
- UK Prospective Diabetes Study (UKPDS) Group: Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet*, 1998, 352, s. 854–855.
- Monnier, L., et al.: The loss of postprandial glycaemic control precedes stepwise deterioration of fasting with worsening diabetes. *Diabetes Care*, 2007, 30, s. 263–269.
- Mittraku, A. – Kelley, D. – Mekan, M., et al.: Role of reduced suppression of glucossem production and diminished early insulin release in impaired glucose tolerance. *N Engl J Med*, 1992, 326, s. 22–29.
- Balvan, P. A. – Podonsky, K. S. – Bowden, D. W., et al.: B-cell failure in type 2 diabetes: postulated mechanisms and prospects for prevention and treatment. *Diabetes Care*, 2014, 37, s. 1751–1758.
- Poitout, V. – Robertson, R. P.: Glucolipotoxicity: fuel excess and  $\beta$ -cell dysfunction. *Endocr Rev*, 2008, 29, s. 351–366.
- Rahier, J. – Guiot, Y. – Goebbels, R. M., et al.: Pancreatic  $\beta$ -cell mass in European subjects with type 2 diabetes. *Diabetes Obes Metab*, 2008, 10 (dopl.), s. 32–42.
- Kramer, C. K. – Zinman, B. – Retnakaran, R.: Short-term intensive insulin therapy in type 2 diabetes mellitus: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol*, 2013, 1, s. 28–34.
- Chen, H. S. – Wu, T. E. – Jap, T. S., et al.: Beneficial effects of insulin on glycaemic control and  $\beta$ -cell function in newly diagnosed type 2 diabetes with severe hyperglycemia after short-term intensive insulin therapy. *Diabetes Care*, 2008, 31, s. 1927–1932.
- Xu, W. – Li, Y. B. – Deng, W. P., et al.: Remission of hyperglycemia following intensive insulin therapy in newly diagnosed type 2 diabetic patients: a long-term follow-up study. *Chin Med J (Engl)*, 2009, 122, s. 2554–2559.
- Pistrosch, F. – Kohler, C. – Schaper, F., et al.: Effects of insulin glargine versus metformin on glycaemic variability, microvascular and  $\beta$ -cell function in early type 2 diabetes. *Acta Diabetol*, 2013, 50, s. 587–595.
- Forst, T. – Larbig, M. – Hohberg, C., et al.: Adding insulin glargine vs. NPH insulin to metformin results in a more efficient postprandial  $\beta$ -cell protection in individuals with type 2 diabetes. *Diabetes Obes Metab*, 2010, 12, s. 437–441.
- Hanefeld, M. – Fleischmann, H. – Schiffhorst, G., et al.: Predictors of response to early basal insulin treatment in patients with type 2 diabetes—the EARLY experience. *Diabetes Technol Ther*, 2014, 16, s. 241–246.
- Fonseca, V. – Gill, J. – Zhou, R., et al.: An analysis of early insulin glargine added to metformin with or without sulfonylurea: impact on glycaemic control and hypoglycaemia. *Diabetes Obes Metab*, 2011, 13, s. 814–822.
- UK Prospective Diabetes Study (UKPDS) Group: Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*, 1998, 352, s. 837–853.
- Buse, J. B. – Peters, A. – Russell-Jones, D., et al.: Is insulin the most effective injectable antihyperglycaemic therapy? *Diabetes Obes Metab*, 2015, 2, s. 145–151.
- ORIGIN Trial Investigators: Characteristics associated with maintenance of mean A1C  $\leq$  6.5% in people with dysglycemia in the ORIGIN trial. *Diabetes Care*, 2013, 36, s. 2915–2922.
- Pennartz, C. – Schenker, N. – Menge, B. A., et al.: Chronic reduction of fasting glycemia with insulin glargine improves first- and second-phase insulin secretion in patients with type 2 diabetes. *Diabetes Care*, 2011, 34, s. 2048–2053.
- Bunck, M. C. – Diamant, M. – Cornér, A., et al.: One-year treatment with exenatide improves  $\beta$ -cell function, compared with insulin glargine, in metformin-treated type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care*, 2009, 32, s. 762–768.
- Hanefeld, M., et al.: Early treatment with basal insulin glargine in people with type 2 diabetes: Lessons from ORIGIN and other cardiovascular trials. *Diabetes Ther*, 2016, 7, s. 187–201.
- Weng, J., et al.: Effect of intensive insulin therapy on beta-cell function and glycaemic control in patients with newly diagnosed type 2 diabetes: a multicentre randomised parallel-group trial. *Lancet*, 2008, 371, s. 1753–1760.
- Gerstein, H. C., et al.: Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med*, 2012, 367, s. 319–328.